

Z.Arbeitsgruppe Öst. Ent.	53	117-120	Wien, 30. 9. 2001	ISSN 0375-5223
---------------------------	----	---------	-------------------	----------------

Description of a new species of the *C. syriacus* group of *Cardiophorus* ESCHSCHOLTZ, 1829 (Coleoptera: Elateridae: Cardiophorinae)

Giuseppe PLATIA & Peter C. CATE

Abstract

Cardiophorus syriacoides sp. n., up to the present confused with *C. syriacus* (Linnaeus, 1758), is described upon specimens from Iran, Georgia, Armenia, Azerbaijan, Turkmenistan and Tadzhikistan. A key to the species of the *C. syriacus* group is from the region given.

Key words: Coleoptera, Elateridae, *Cardiophorus*, new species, Iran, Georgia, Armenia, Azerbaijan, Turkmenistan, Tadzhikistan.

Zusammenfassung

Cardiophorus syriacoides sp. n., bisher mit *C. syriacus* (Linnaeus, 1758) vermischt, wird nach Exemplaren aus Iran, Georgien, Armenien, Azerbaijan, Turkmenistan und Tadzhikistan beschrieben. Ein angeschlossener Bestimmungsschlüssel dient der Determination von Arten der *C. syriacus* Gruppe dieses Gebietes.

Introduction

During studies on click-beetle species from Iran our attention was attracted to specimens determined as *Cardiophorus syriacus* (Linnaeus, 1758), which proved upon closer examination of the morphological features to differ markedly from specimens standing under this name from Syria, the type locality, and Turkey. The examination of specimens from Iran, Georgia, Armenia, Azerbaijan, Turkmenistan and Tadzhikistan show that many also belong to the new species. *C. syriacoides* sp. n. can be recognized by the longer antennae in male and female, by the sparser punctuation of pronotum and by a different structure of male and female genitalia.

The *Cardiophorus syriacus* group contains four species in the region under study. *C. hinkei* Frivaldszky, 1837, is reported from Bulgaria, Greece and western Turkey (PLATIA & GUDENZI, 2000a). *C. syriacus* was originally described from Syria and is also reported from Turkey, Armenia, Azerbaijan, Iran and Turkmenistan (CANDÈZE, 1860; JAKOBSON, 1913; SCHENKLING, 1925; GURJEVA, 1966; MARDJANIAN, 1977, 1987; AGAEV, 1988; DOLIN & ATAMURADOV, 1994; PLATIA & GUDENZI, 2000b). Specimens from Turkey and probably also those from Armenia (according to drawings of the female genital in MARDJANIAN (1977, 1987)) belong to *C. syriacus*, whereas those mentioned in GURJEVA (1966) and DOLIN & ATAMURADOV (1994) from Turkmenistan as well as former records from Iran are very probably *C. syriacoides* sp. n. AGAEV (1985) described *C. ordubadensis* from Azerbaijan, which he distinguishes from *C. syriacus* by a uniform body colour, longer

antennae and the male and female genitalia. We have examined 5 specimens of this species, four from Azerbaijan and one from Erivan, Armenia (new country record).

As the new species is also known from Georgia, Armenia and Azerbaijan, the ranges of *C. syriacus* and *C. syriacoides* could very well overlap in the Caucasus region. Furthermore, as we have seen a specimen of *C. hinkei* labelled Elisabethpol (Azerbaijan, new country record), the ranges of *C. hinkei* and *C. syriacoides* apparently do overlap. Only further studies on specimens from this area can elucidate the distribution of all four species.

***Cardiophorus syriacoides* sp. n. (Photos 1, 2, 3)**

Material examined: holotype ♀, **Iran**: Ala-Dagh, Budschmurd, 1033m, vi.1902, ex. Hauser coll. (Natural History Museum, Vienna); 31 paratypes (4 ♀ ♀, 10 ♂ ♂; 17 not sexed), Astrabad, v.1899, ex Hauser coll.; **Azerbaijan**: Elisabethpol, ex Kolenati coll.; Talysch, Larik distr., env. Gosmaljan, leg. S. Kasantsev, 6.vi.1988; **Armenia**: Megri, Armenia, Viktorov & Dlusskij, 17.iv.1959; Armenia, Megri, 30.iv.1964; **GEORGIA**: GSSR, Vaschlavan conservation area, 15.v.1981, V. G. Dolin (ex larva); **Turkmenistan**: Transkaspia, Saramsakli, vi.1907, ex Hauser coll.; Kov-Ata, Ibachardenli, 110 km NW Aschabad, 10.v.1989, leg. Wrase; Firjuza/Kopet Dag, 22.iv.1981, leg. J. Strnad; Firjuza, 26.v.1983, Kh. Atamuradov; Kara-Kale, Ak-Dere gorge, 1500m, 25.v.1977, leg. W. Dolin; Transcaspi, Tschandyl, AK, 29.iv.1933; Transcaspi, Tschandyl, Kara-K., 25.iv.1933; Transcaspi, Kara-Kala, 20.iv.1933, Ut.; Turkmenistan, Z. Kopetdag, 720m, m. Taldycharshma, 24.iv.1989, D.W.; Turkmenistan, slopes nr. Kizil-Arbat, 15.iv.1975; hills and slopes of Kopet-Dagh, 12 km W Kizil-Arbat, 12.iv.1953; Turkmenistan, Tschuli gorge, slopes, on Crataegus, 3.v.1975; Turkmenistan, Kara-Kala reg., Tschendyr gorge, 20.v.1975, leg. V. Dolin; Turkmenistan, Umg. Aschgabat, Kalininsk, 15.iv.1976, V. G. Dolin; **Tadzhikistan**: Bulbulasma, 18.iv.1980, leg. J. Strnad; **Unknown locality**: 2 km S Megrenjank(?), 18.iv.1987. (Natural History Museum, Vienna; coll. Cate, Vienna; coll. Platia, Gatteo; Schmalhausen Institute of Zoology, Kiev).

Description of female: Bicoloured; pronotum including propleurae red-orange; head, scutellum, elytra, prosternum, thorax and abdomen (sometimes last two sternites reddish) black; antennae blackish with base of segments ferruginous; legs ferruginous, generally with darkened femora. Elytra with whitish pubescence forming two distinctive transversal bands at base and in posterior half.


Frons flat to moderately convex, impressed only at anterior margin, the latter regularly arcuate and feebly thickened; punctures broad and contiguous. Antennae reaching tips of the posterior angles of pronotum; second article subconical, shorter than third; third to tenth subtriangular, approximately the same length; eleventh subellipsoidal, pointed at apex.

Pronotum as long as wide, strongly and regularly convex with a vestige of median longitudinal furrow in the basal declivity; sides strongly arcuate with the greatest width in the middle; posterior angles short, truncate, not divergent, briefly carinate; lateral suture complete; punctuation uniform, on the disk with punctures of approximately the same size, simple, with shiny and very small intervals.

Scutellum heart-shaped, as long as wide, slightly concave and emarginate in the middle of base, densely punctured. Elytra on average 2.3 times longer than pronotum and twice as long as wide; convex on the disk, suboval; striae well-marked and punctured; interstriae flat, very densely and finely punctured.

Claws simple.

Bursa copulatrix (photo 2) with two symmetrical sclerified plates and a median U-shaped sclerite.



Cardiophorus syriacoides sp. n.: Photo 1: Aedeagus in dorsal view (paratype) (length: 0.92 mm). Photo 2: Sclerifications in the bursa copulatrix (holotype) (length of plate: 0.47 mm; length of median piece: 0.4 mm). Photo 3: Adult male (paratype) (length: 5.1 mm).

Male (photo 3): Similar to female but generally with a more slender body and longer antennae exceeding the tips of posterior angles of pronotum by one to two segments.

Aedeagus (photo 1) with parameres simple at apex.

Length: 5.1-7.5 mm; width: 1.6-2.3 mm.

Key to species of the *Cardiophorus syriacus* group

1. First two antennomeres reddish..... 2
- First two antennomeres blackish..... *syriacus* (Linnaeus)
2. Antennae shorter, reaching (male) or not reaching (female) tips of posterior angles of pronotum; pronotal punctures denser, contiguous..... *hinkei* Frivaldszky
- Antennae longer, exceeding (male) or reaching (female) tips of posterior angles of pronotum; pronotal punctures sparser, not contiguous..... 3
3. Body black with red-orange pronotum and propleurae, antennae exceeding tips of posterior angles of pronotum by 1-2 articles (male)..... *syriacoides* sp. n.
- Body generally red-brown with pronotum and propleura of same colour as elytra or only slightly lighter (darker specimens also have darkened pronotum), antennae exceeding tips of posterior angles of pronotum by 2-3 articles (male)..... *ordubadensis* Agaev

Besides the characters mentioned in the key, the examination of male and female genitalia are of great value in distinguishing the species.

Acknowledgements

We want to thank Dr. H. Schönmann of the Natural History Museum of Vienna for allowing us to study specimens of the new species in his care. We also thank Prof. V. G. Dolin, Schmalhausen Institute of Zoology, Kiev, for being able to examine further specimens.

LITERATURE

- AGAEV, B. I. 1985: Novye vidy zhulkov-shchelkunov (Coleoptera, Elateridae) iz vostochnogo Zakavkasya. [New species of click-beetles (Col. Elateridae) from Eastern Transcaucasia.] - Vestnik Zoologii, 1985(6):56-58.
- AGAEV, B. I. 1988: Zhestkokryye shchelkyny v biotsenozach Azerbaydzhana. [The click-beetles of the biocoenoses of Azerbaijan.] - Azerbaydzhanskoe gosudarstvennoe izdatel'stvo. Baku. 1988. 120 pp.
- CANDÈZE, E. C. A. 1860: Monographie des Élatérides. III. - Mémoires de la Société Royale des Sciences de Liège, 15: 512 pp., 5 pls.
- DOLIN V. G. & ATAMURADOV, Kh. I. 1994: Zhuki-shchelkuny (Elateridae) Turkmenistana. [Click-beetles (Elateridae) of Turkmenistan.] - Institut zoologii NAN Ukrainy. Kiev. 1994. 178 pp.
- GURJEVA, E. L. 1966: Zhuki-shchelkunovy podsemeystva Cardiophorinae (Coleoptera, Elateridae) sredney Asii. In: Novye vidy nasekomych fauny SSSR i sopredel'nykh stran. [Click-beetles of the subfamily Cardiophorinae (Coleoptera, Elateridae) of Middle Asia. In: New species of insects of the fauna of the USSR and neighbouring territories] - Trudy Zoologicheskogo Instituta Akademii Nauk SSSR, 37: 62-97.
- JAKOBSON G. G. 1913: Zhuki Rossii i Zapadnoj Evropy. Rukovodstvo k' opredeleniyu zhukov'. [The Beetles of Russia and Western Europe. A handbook for the determination of beetles.] Vol. 10. - A. F. Devrien. St. Petersburg. 1024 pp.
- MARDJANIAN, M. A. 1977: Izmenchivost' khitinizirovannikh struktur genitaliy shchelkunov roda Cardiophorus (Coleoptera, Elateridae). [Variation of chitinized structures of the genitalia in click beetles of the genus Cardiophorus Eschz. (Coleoptera, Elateridae).] - Zoologicheskiy Zhurnal, 56(11): 1629-1636.
- MARDJANIAN, M. A. 1987: Fauna Armyanskoi SSR. Nasekomye Zhestkokryle. Shchelkuny (Elateridae). - Izdatelstvo Akademii Nauk Armanskoi SSR. Erevan. 204 pp.
- PLATIA, G. & GUDENZI, I. 2000a: Contributo alla conoscenza dei *Cardiophorus* della Grecia. (Coleoptera, Elateridae). - Bollettino della Società entomologica italiana, Genova, 132(82): 135-155.
- PLATIA, G. & GUDENZI, I. 2000b: Revisione dei *Cardiophorus* Eschscholtz della Turchia. I: Le specie bicolori. (Coleoptera, Elateridae, Cardiophorinae). Lambillionea, 100(4): 601-612.
- SCHENKLING, S. 1925: Coleopterorum Catalogus auspiciis et auxilio W. Junk. Pars 80, Elateridae I. - W. Junk. Berlin. 263 pp.

Addresses of the authors: Dr. PLATIA Giuseppe, Via Molino Vecchio, 21, 47030 Gatteo (FO), Italy:
e-mail: pinoplatia@libero.it

Dr. Peter C. CATE, Bundesamt und Forschungszentrum für Landwirtschaft,
Spargelfeldstr. 191, A-1226 Vienna, Austria; e-mail: peter.cate@bfl.ac.at

ZOBODAT - www.zobodat.at

Zoologisch-Botanische Datenbank/Zoological-Botanical Database

Digitale Literatur/Digital Literature

Zeitschrift/Journal: [Zeitschrift der Arbeitsgemeinschaft Österreichischer Entomologen](#)

Jahr/Year: 2001

Band/Volume: [53](#)

Autor(en)/Author(s): Platia Giuseppe, Cate Peter Carl

Artikel/Article: [Description of a new species of the *C. syriacus* group of Cardiophorus Eschscholtz, 1829 \(Coleoptera: Elateridae: Cardiophorinae\).](#)
[117-120](#)